To appear in the *Journal of Nonparametric Statistics*
Vol. 00, No. 00, Month 20XX, 1–26

arXiv:1410.2507v2 [math.PR] 20 Feb 2015

Gamma kernel estimation of multivariate density and its derivative on the nonnegative semi-axis by dependent data

L. Markovich[a*]

(*Received 00 Month 20XX; in final form 00 Month 20XX*)

We consider the nonparametric estimation of the multivariate probability density function and its partial derivative with a support on $[0,\infty)$ by dependent data. To this end we use the class of kernel estimators with asymmetric gamma kernel functions. The gamma kernels are nonnegative. They change their shape depending on the position on the semi-axis and possess good boundary properties for a wide class of densities. The theoretical asymptotic properties of the multivariate density and its partial derivative estimates like biases, variances and covariances are derived. We obtain the optimal bandwidth selection for both estimates as a minimum of the mean integrated squared error (MISE) assuming dependent data with a strong mixing. Optimal rates of convergence of the MISE are found.

Keywords: density derivative; multivariate dependent data; gamma kernel; nonparametric estimation

AMS Subject Classification: 62G07; 62G20; 62H12

1. Introduction

Nonnegatively supported probability density functions (pdf) can be used to model a wide range of applications in engineering, signal processing, medical research, quality control, actuarial science and climatology among others. In the optimal filtering in the signal processing and control of nonlinear processes the exponential pdf class is used, (**?**). Most total insurance claim distributions are shaped like gamma pdfs (**?**). The gamma distribution are also used to model rainfalls (**?**). Erlang and χ^2 pdfs are widely used in modeling insurance portfolios (**?**).

The wide use of these pdfs in practice leads to need of their estimation by finite data samples. One of the most common nonparametric pdf estimation methods are kernel estimators. The kernel density estimators were originally introduced for univariate independent identically distributed (iid) data in (**??**) for symmetrical kernels. The latter approach was widely adopted for tasks where a support of the underlying pdf f is unbounded. In case when the pdf has a nonsymmetric support $[0,\infty)$ the problem of the large bias on the zero boundary appears. This leads to a bad quality of the estimates (**?**). Obviously, the boundary bias for the multivariate pdf estimation becomes even more solid (**?**). To overcome this problem one can use special approaches such as the data reflection (**?**), the boundary kernels (**?**), the hybrid method (**?**), the local linear estimator (**??**) among others. Another solution is to use the asymmetrical kernel functions

*Corresponding author. Email: kimo1@mail.ru [a]*Institute of Control Sciences, Russian Academy of Sciences, Moscow, Russia.*

instead of symmetrical ones. For univariate nonnegative iid random variables (r.v.s), the estimators with gamma kernels were proposed in (**?**). The gamma kernel estimator was developed for univariate dependent data in (**?**). In (**?**) the gamma kernel estimator of the multivariate pdf for the nonnegative iid r.v.s was introduced.

The gamma kernel is nonnegative and flexible regarding the shape. This allows to fit accurate the multi-modal pdfs and their derivatives. Estimators constructed with gamma kernels have no boundary bias if $f''(0) = 0$ holds, i.e when the underlying density $f(x)$ has a shoulder at $x = 0$ (see formula (4.3) in (**?**)). This shoulder property is fulfilled, for example, for a wide exponential class of pdfs. Other asymmetrical kernel estimators like inverse Gaussian and reciprocal inverse Gaussian estimators were studied in (**?**). The comparison of these asymmetric kernels with the gamma kernel is given in (**?**).

Along with the density estimation it is often necessary to estimate the derivative of the pdf. The estimation of the univariate density derivative by the gamma kernel estimator was proposed by authors in (**??**) for iid data and in (**?**) for a strong mixing dependent data. Our procedure achieves the optimal MISE of order $n^{-\frac{4}{7}}$ when the optimal bandwidth is of order $n^{-\frac{2}{7}}$. In (**?**) an optimal MISE of the kernel estimate of the first derivative of order $n^{-\frac{4}{7}}$ corresponding to the optimal bandwidth of order $n^{-\frac{1}{7}}$ for symmetrical kernels was indicated.

In this paper, we introduce the gamma product kernel estimators for the multivariate density with the nonnegative support and its partial derivative by the multivariate dependent data with a strong mixing. The asymptotical behavior of the estimates and the optimal bandwidths in the sense of minimal MISE are obtained. The derivative estimation requires a specific bandwidth different from that for the pdf estimation.

The pdf derivative could be useful to find the slope of the pdf curve, its local extremes, saddle points, significant features in data as well as in regression analysis (**?**). The pdf derivative also plays a key role in clustering via mode seeking (**?**). Furthermore, we are able to write down the estimator for the logarithmic derivative of the nonnegative pdf for the multivariate dependent data. The estimate of the logarithmic derivative of the pdf is necessary for an optimal filtering in the signal processing and control of nonlinear processes (**?**).

The outline of the paper is as follows. In Section 2 we define the gamma kernel estimators of the multivariate density and its partial derivative. In Section 3 we obtain the bias, the variance and the covariance of the pdf estimate. Using these results we derive the optimal bandwidth and the corresponding rate of the optimal MISE. In Section 4 we obtain the same for the estimate of the partial pdf derivative.

2. Gamma Kernel Estimation

Let $\{\mathbf{X}_i = (X_{i1}, \dots, X_{id})^T\}_{i=1}^n$ be a strongly stationary sequence of d-dimensional variables with an unknown probability density function $f(x_1^d)$, $x_1^d = (x_1, \dots, x_d)^T$ which is defined on the bounded support $x_1^d \in \mathbf{R}^{+d}$. We assume that the sequence $\{\mathbf{X}_s\}$ is $\alpha-$mixing with coefficient

$$\alpha(k) = \sup_t \sup_{\substack{A\in\sigma\{X_s, s\le t\} \\ B\in\sigma\{X_s, s\ge t+k\}}} |P(A\cap B) - P(A)P(B)|, \quad k \ge 1.$$

Here $\alpha(k) \to 0$ as $k \to \infty$. The α-mixing condition, also called strong mixing, is satisfied by many stochastic processes, i.e. autoregressive process (**?**). For these sequences we will use a notation $\{\mathbf{X}_s\} \in \mathcal{S}(\alpha)$. To estimate the unknown pdf we use the product gamma

kernel estimator (see for example in (**?**))

$$\widehat{f}(x_1^d) = \frac{1}{n}\sum_{i=1}^{n}\prod_{j=1}^{d} K_{\rho(x_j.b_j),b_j}\left(X_{ij}\right), \tag{1}$$

where $b \equiv \{b_j\}_{j=1}^{d}$ is the vector of the bandwidth parameters such that $b \to 0$ as $n \to \infty$. As a kernel functions for each variable we propose the gamma kernel (**?**)

$$K_{\rho(x,b),b}(t) = \frac{t^{\rho(x,b)-1}\exp(-t/b)}{b^{\rho(x,b)}\Gamma(\rho(x,b))}, \tag{2}$$

where $\Gamma(\cdot)$ is a standard gamma function. To improve the properties of the gamma kernel estimator (**?**) used parameter $\rho(x,b)$ defined as

$$\rho(x,b) = \begin{cases} \rho_1(x,b) = x/b, & \text{if} \quad x \geq 2b, \\ \rho_2(x,b) = (x/(2b))^2 + 1, & \text{if} \quad x \in [0,2b). \end{cases} \tag{3}$$

Since gamma kernel is nonnegative it has no weight on the negative semiaxes in contrast to symmetrical kernel estimators. Hence, the use of the gamma kernels is more natural for estimation of the nonnegative supported pdfs.

In nonparametric filtration of stochastic signals the single series of realizations x_1^n is divided into the fragments $x_k^{k+\tau}$, $1 \leq k \leq n-\tau-1$. The last fragment $x_{n-\tau}^n$ is used as the argument of the pdf, parameter $0 \leq \tau \leq n-1$ is called the degree of dependence (**?**). Hence, without loss of generality we shall estimate the pdf $f(x_{n-\tau}^n)$. Its kernel estimate is

$$\widehat{f}(x_{n-\tau}^n) = \frac{1}{n}\sum_{i=1}^{n}\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j),b_j}\left(X_{ij}\right). \tag{4}$$

Remark 1 If $\tau = n-1$ the latter estimate is nothing but (1) where $d = n$.

Using (3) the pdf estimator (24) can be written as

$$\widehat{f}(x_{n-\tau}^n) = \begin{cases} \frac{1}{n}\sum\limits_{i=1}^{n}\prod\limits_{j=n-\tau}^{n} K_{\rho_1(x_j,b_j),b_j}\left(X_{ij}\right), \text{if} \quad x \geq 2b, \\ \frac{1}{n}\sum\limits_{i=1}^{n}\prod\limits_{j=n-\tau}^{n} K_{\rho_2(x_j,b_j),b_j}\left(X_{ij}\right), \text{if} \quad x \in [0,2b). \end{cases} \tag{5}$$

In (**??**) the derivative of the univariate pdf was estimated just like the derivative of the gamma kernel estimator. For the multivariate case, we can analogically estimate any partial derivative of $f(x_{n-\tau}^n)$. For example, the partial derivative of $f(x_{n-\tau}^n)$ by x_n can be estimated as

$$\widehat{f}'_{x_n}(x_{n-\tau}^n) = \frac{1}{n}\sum_{i=1}^{n}\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j),b_j}\left(X_{ij}\right)\left(K_{\rho(x_n,b_n),b_n}\left(X_{in}\right)\right)'_{x_n}, \tag{6}$$

where

$$\left(K_{\rho(x,b),b}(t)\right)'_{x_n} = \begin{cases} K'_{\rho_1(x,b),b}(t) = \frac{1}{b}K_{\rho_1(x,b),b}(t)L_1(t,x,b), & \text{if} \quad x \geq 2b, \\ K'_{\rho_2(x,b),b}(t) = \frac{x}{2b^2}K_{\rho_2(x,b),b}(t)L_2(t,x,b), & \text{if} \quad x \in [0,2b), \end{cases} \tag{7}$$

is the partial derivative of (2) and

$$L_i(t,x,b) = \ln t - \ln b - \Psi(\rho_i(x,b)), \quad i = 1,2. \tag{8}$$

Here, $\Psi(\rho_i(x,b))$ denotes the Digamma function that is the logarithmic derivative of the gamma function.

Remark 2 The mathematical technic applied for the derivative estimation is similar to one applied for the pdf. However, formulas became much more complicated particulary because one has to deal with the special Digamma function that includes the bandwidth vector b. Hence, one has to pick out the order by b from complicated expressions containing logarithms and the special function.

Using (7), formula (6) can be rewritten as

$$\widehat{f}'_{x_n}(x^n_{n-\tau}) = \begin{cases} \frac{1}{n}\sum\limits_{i=1}^{n} \frac{1}{b_n} L_1(X_{in}, x_n, b_n) \prod\limits_{j=n-\tau}^{n} K_{\rho_1(x_j,b_j),b_j}(X_{ij}), & \text{if} \quad x \geq 2b, \\ \frac{1}{n}\sum\limits_{i=1}^{n} \frac{x_n}{2b_n^2} L_2(X_{in}, x_n, b_n) \prod\limits_{j=n-\tau}^{n} K_{\rho_2(x_j,b_j),b_j}(X_{ij}), & \text{if} \quad x \in [0, 2b). \end{cases} \tag{9}$$

As the measure of error of the proposed estimators (5) and (9) it is natural to consider the MISEs which are defined as

$$MISE(\widehat{f}(x^n_{n-\tau})) = \mathsf{E}\int_0^\infty (f(x^n_{n-\tau}) - \widehat{f}(x^n_{n-\tau}))^2 dx, \tag{10}$$
$$MISE(\widehat{f}'_{x_n}(x^n_{n-\tau})) = \mathsf{E}\int_0^\infty (f'_{x_n}(x^n_{n-\tau}) - \widehat{f}'_{x_n}(x^n_{n-\tau}))^2 dx.$$

The unknown smoothing parameters of (5) and (9) are obtained as the minima of (10).

Remark 3 The integrals in (10) can be splitted into two integrals $\int_0^{2b}$ and $\int_{2b}^\infty$. Further we shall do all the proofs for the case when $x \geq 2b$ because the integral $\int_0^{2b}$ tends to zero when $b \to 0$. Hence, we omit the indices for $\rho_1(x_j, b_j)$, $L_1(X_{ik}, x_k, b_k)$ and use instead $\rho(x_j, b_j)$, $L(X_{ik}, x_k, b_k)$.

It is known that the mean squared error (MSE) for dependent r.v.s is determined as

$$MSE(\widehat{f}(x^n_{n-\tau})) = (Bias(\widehat{f}(x^n_{n-\tau}))^2 + Var(\widehat{f}(x^n_{n-\tau})), \tag{11}$$
$$MSE(\widehat{f}'_{x_n}(x^n_{n-\tau})) = (Bias(\widehat{f}'_{x_n}(x^n_{n-\tau}))^2 + Var(\widehat{f}'_{x_n}(x^n_{n-\tau})),$$

where the variances contain the covariance terms

$$Var(\widehat{f}(x_{n-\tau}^n)) = Var\left(\frac{1}{n}\sum_{i=1}^{n}\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j),b_j}(X_{ij})\right) \tag{12}$$

$$= \frac{1}{n^2}\left(\sum_{i=1}^{n} Var\left(\widetilde{\widetilde{K}}(X_i,x,b)\right) + \sum_{i,k=1,i\neq k}^{n} Cov\left(\widetilde{\widetilde{K}}(X_i,x,b),\widetilde{\widetilde{K}}(X_k,x,b)\right)\right)$$

$$= \frac{1}{n}var(\widehat{f}(x_{n-\tau}^n)) + \frac{2}{n}\sum_{i=1}^{n}\left(1-\frac{i}{n}\right)Cov\left(\widetilde{\widetilde{K}}(X_i,x,b),\widetilde{\widetilde{K}}(X_{1+i},x,b)\right),$$

$$Var(\widehat{f}'_{x_n}(x_{n-\tau}^n)) = Var\left(\frac{1}{n}\sum_{i=1}^{n}\frac{1}{b_n}L(X_{in},x_n,b_n)\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j),b_j}(X_{ij})\right) \tag{13}$$

$$= \frac{1}{n^2b_n^2}\left(\sum_{i=1}^{n} Var\left(\widetilde{K}(X_i,x,b)\right) + \sum_{i,k=1,i\neq k}^{n} Cov\left(\widetilde{K}(X_i,x,b),\widetilde{K}(X_k,x,b)\right)\right)$$

$$= \frac{1}{nb_n^2}var(\widehat{f}'_{x_n}(x_{n-\tau}^n)) + \frac{2}{nb_n^2}\sum_{i=1}^{n}\left(1-\frac{i}{n}\right)Cov\left(\widetilde{K}(X_1,x,b),\widetilde{K}(X_{1+i},x,b)\right).$$

Here and after we use the following notations

$$\widetilde{\widetilde{K}}(X_i,x,b) = \prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j),b_j}(X_{ij}), \quad \widetilde{K}(X_i,x,b) = L(X_{in},x_n,b_n)\widetilde{\widetilde{K}}(X_i,x,b).$$

3. Convergence rate of the density estimator

In this section we obtain the asymptotic properties of the estimator (5). To this end we derive the bias, the variance and the covariance determined in (11) in the following lemmas. All proofs hold assuming that all the components of the bandwidths vector are different. For the practical use we give the simpler formulation of lemmas for the equal bandwidths $b_1 = b_2 = \ldots = b_n = b$.

LEMMA 3.1 *If $b_1 = b_2 = \ldots = b_n = b$ and $b \to 0$ as $n \to \infty$, then the bias for the pdf estimate* (5) *is equal to*

$$Bias(\widehat{f}(x_{n-\tau}^n)) = \frac{b}{2}\sum_{j=n-\tau}^{n} x_j\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} + o(b). \tag{14}$$

LEMMA 3.2 *If $b_1 = b_2 = \ldots = b_n = b$ and $b \to 0$, $nb^{\frac{\tau+1}{2}} \to \infty$ as $n \to \infty$, then the*

variance expansion for the pdf estimate (5) *is equal to*

$$var(\widehat{f}(x_{n-\tau}^n)) = \frac{b^{-\frac{(\tau+1)}{2}}}{n}\left(\prod_{j=n-\tau}^{n}\frac{x_j^{-1/2}}{2\sqrt{\pi}}\right)\left(f(x_{n-\tau}^n)+bv_1(x_{n-\tau}^n)+b^2v_2(x_{n-\tau}^n)\right)$$
$$-\frac{1}{n}\left(f(x_{n-\tau}^n)+\frac{1}{2}\sum_{j=n-\tau}^{n}x_jb\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2}\right)^2+o(b^2), \tag{15}$$

where

$$v_1(x_{n-\tau}^n) = \sum_{j=n-\tau}^{n}\left(-\frac{1}{2}\frac{\partial f(x_{n-\tau}^n)}{\partial x_j}+\frac{x_j}{4}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2}\right),$$
$$v_2(x_{n-\tau}^n) = -\sum_{j=n-\tau}^{n}\sum_{i=n-\tau}^{n}\frac{x_j}{8}\frac{\partial^3 f(x_{n-\tau}^n)}{\partial x_j^2\partial x_i}. \tag{16}$$

LEMMA 3.3 *Let*

(1) $\{X_j\}_{j\geq 1}\in\mathcal{S}(\alpha)$ *and* $\int_1^\infty \alpha(\tau)^\upsilon d\tau<\infty,\quad 0<\upsilon<1$ *hold,*
(2) $f(x_{n-\tau}^n)$ *be a twice continuously differentiable function,*
(3) $b\to 0$ *and* $nb^{(\tau+1)\frac{\upsilon+1}{2}}\to\infty$ *as* $n\to\infty$.

Then, the covariance is bounded by

$$|C(\widehat{f}(x_{n-\tau}^n)| \leq \frac{b^{-(\tau+1)\frac{\upsilon+1}{2}}}{n}\int_1^\infty \alpha(\tau)^\upsilon d\tau$$
$$\cdot D(\upsilon,x_{n-\tau}^n)\left(bS(\upsilon,x_{n-\tau}^n)+f(x_{n-\tau}^n)\frac{3\upsilon-1}{2(\upsilon-1)}\right)^{1-\upsilon}+o\left(b^2\right), \tag{17}$$

where

$$S(\upsilon,x_{n-\tau}^n) = \sum_{i=n-\tau}^{n}\frac{\upsilon+1}{(\upsilon-1)^2x_i}f(x_{n-\tau}^n)+\frac{\upsilon+1}{\upsilon-1}\frac{\partial f(x_{n-\tau}^n)}{\partial x_i}+\frac{x_i}{2}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_i^2},$$

$$D(\upsilon,x_{n-\tau}^n) = 2(2\pi)^{-\frac{\tau(\upsilon+1)+\upsilon-1}{2}}\left(\prod_{j=n-\tau}^{n}x_j^{-\frac{\upsilon+1}{2}}\right).$$

The proofs of Lemmas 3.1 - 3.3 are given in Appendices .1, .2, .3. Using the results of the latter the MISE (10) can be written. Hence, the following theorem holds.

THEOREM 3.4 *In the conditions of Lemmas 3.1 - 3.3, the optimal bandwidth that pro-*

vides minimum of the MISE is

$$b=\left(\frac{(\tau+1)(\upsilon+1)(3\upsilon-1)^{1-\upsilon}}{n(2\upsilon-2)^{1-\upsilon}}\frac{\int D(\upsilon,x_{n-\tau}^{n})f(x_{n-\tau}^{n})^{1-\upsilon}dx_{n-\tau}^{n}}{\int\left(\sum\limits_{j=n-\tau}^{n}x_{j}\frac{\partial f(x_{n-\tau}^{n})}{\partial x_{j}^{2}}\right)^{2}dx_{n-\tau}^{n}}\int\limits_{1}^{\infty}\alpha(\tau)^{\upsilon}d\tau\right)^{\frac{\tau(\upsilon+1)+\upsilon+5}{2}} \tag{18}$$

where we denote multiple integrals as $\int\ldots\int dx_{n-\tau}\ldots dx_{n}\equiv\int dx_{n-\tau}^{n}$.

The proof is given in Appendix .4.

Obviously, it is impossible to use the bandwidth (18) since it contains the unknown mixing coefficient $\alpha(\tau)$. The following lemma serves to overcome this problem.

LEMMA 3.5 *Let the conditions of Lemma 3.3 be satisfied. If additionally there exist* $M>0$ *such that for all* $\tau>1$ *and* $(x,y)\in\mathbb{R}^{2}$ *it follows*

$$|f_{\tau}(x,y)-f(x)f(y)|\leq M,$$

then the covariance of the multivariate pdf estimator is

$$|C(\widehat{f}(x_{n-\tau}^{n})|\leq\frac{2}{n}\sum_{\tau=2}^{c(n)}Cov\left(\widetilde{\widetilde{K}}\left(X_{1},x,b\right),\widetilde{\widetilde{K}}\left(X_{\tau},x,b\right)\right)$$
$$+\frac{2}{n}\sum_{\tau=c(n)+1}^{\infty}Cov\left(\widetilde{\widetilde{K}}\left(X_{1},x,b\right),\widetilde{\widetilde{K}}\left(X_{\tau},x,b\right)\right)|=I_{1}+I_{2}$$

and its rate of convergence is

$$I_{1}\leq\frac{2M}{nb^{\frac{\tau+1}{8}}}, \tag{19}$$
$$I_{2}\leq\frac{D(\kappa,x_{n-\tau}^{n})}{nb^{\frac{(\tau+1)}{16}}}\left(f(x_{n-\tau}^{n})\frac{6\kappa-1}{2(2\kappa-1)}+bS(\kappa,x_{n-\tau}^{n})\right)^{1-2\kappa}\sum_{\tau=2}^{\infty}\tau\alpha(\tau)^{2\kappa}+o\left(b^{2}\right).$$

The proof is given in Appendix .5.

From Lemma 3.5 the following order of the covariance $C(\widehat{f}_{n}(x_{n-\tau}^{n}))\sim n^{-1}b^{-\frac{\tau+1}{8}}$ holds. Thus, the covariance can be neglected in comparison with the variance $var(\widehat{f}_{n}^{\prime}(x_{n-\tau}^{n}))\sim n^{-1}b^{-\frac{\tau+1}{2}}$.

THEOREM 3.6 *In conditions of Lemmas 3.1, 3.2, 3.5, the optimal bandwidth which provides minimum of the MISE is*

$$b=\left(\frac{(\tau+1)\int\limits_{0}^{\infty}\left(\prod\limits_{j=n-\tau}^{n}\frac{x_{j}^{-1/2}}{2\sqrt{\pi}}\right)f(x_{n-\tau}^{n})dx_{n-\tau}^{n}}{\int\limits_{0}^{\infty}\left(\sum\limits_{j=n-\tau}^{n}x_{j}\frac{\partial^{2}f(x_{n-\tau}^{n})}{\partial x_{j}^{2}}\right)^{2}dx_{n-\tau}^{n}}\right)^{\frac{2}{5+\tau}}n^{-\frac{2}{5+\tau}}. \tag{20}$$

The proof is given in Appendix .6.

COROLLARY 3.7 *The result of Theorem 3.6 coincides with the results of (?) for univariate pdf and iid case.*

4. Convergence rate of the density derivative estimator

Similarly to Section 3 we obtain the asymptotic properties of the density derivative estimator (9). The bias, the variance and the covariance determined in (11) are derived in the following lemmas.

LEMMA 4.1 *If $b_1 = b_2 = \ldots = b_n = b$ and $b \to 0$ as $n \to \infty$, then the bias for the density derivative estimate* (9) *is equal to*

$$Bias(\widehat{f}'_{x_n}(x^n_{n-\tau})) = bB_1(x^n_{n-\tau}) + b^2 B_2(x^n_{n-\tau}) + o\left(b\right), \tag{21}$$

where we denote

$$B_1(x^n_{n-\tau}) = \frac{f(x^n_{n-\tau})}{12x_n^2} + \frac{1}{4x_n}\sum_{j=n-\tau}^{n} x_j \frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2},$$

$$B_2(x^n_{n-\tau}) = \frac{1}{24x_n^2}\sum_{j=n-\tau}^{n} x_j \frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2}.$$

LEMMA 4.2 *If $b_1 = b_2 = \ldots = b_n = b$ and $b \to 0$, $nb^{\frac{\tau+3}{2}} \to \infty$ as $n \to \infty$, then the variance of the density derivative estimate* (9) *is equal to*

$$\begin{aligned} var(\widehat{f}'_{x_n}(x^n_{n-\tau})) = \frac{b^{-\frac{(\tau+1)}{2}}}{n}\left(\prod_{j=n-\tau}^{n}\frac{x_j^{-1/2}}{2\sqrt{\pi}}\right)\Bigg(bV_1(x^n_{n-\tau}) + b^2 V_2(x^n_{n-\tau}) \\ + \frac{1}{b}V_3(x^n_{n-\tau}) + V_4(x^n_{n-\tau})\Bigg) - \frac{1}{n}\Bigg(b^2 B_1^2(x^n_{n-\tau}) + \left(\frac{\partial f(x^n_{n-\tau})}{\partial x_n}\right)^2 \\ + 2\frac{\partial f(x^n_{n-\tau})}{\partial x_n}\left(bB_1(x^n_{n-\tau}) + b^2 B_2(x^n_{n-\tau})\right)\Bigg) + o\left(b^2\right), \end{aligned} \tag{22}$$

where

$$V_1(x^n_{n-\tau}) = -\frac{1}{24x_n^2}\frac{\partial f(x^n_{n-\tau})}{\partial x_n} + \sum_{j=n-\tau}^{n}\left(\frac{1}{8x_n}\frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2} - \frac{1}{8x_n^2}\frac{\partial f(x^n_{n-\tau})}{\partial x_j}\right) + \frac{7}{48x_n^3}f(x^n_{n-\tau}),$$

$$V_2(x^n_{n-\tau}) = \frac{7}{576x_n^4}f(x^n_{n-\tau}) + \sum_{j=n-\tau}^{n}\left(\frac{1}{16x_n^2}\frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2} - \frac{7}{96x_n^3}\frac{\partial f(x^n_{n-\tau})}{\partial x_j} + \frac{1}{48x_n^2}\frac{\partial^2 f(x^n_{n-\tau})}{\partial x_n \partial x_j}\right),$$

$$V_3(x^n_{n-\tau}) = \frac{f(x^n_{n-\tau})}{2x_n}, \quad V_4(x) = \frac{f(x^n_{n-\tau})}{4x_n^2} - \sum_{j=n-\tau}^{n} \frac{1}{4x_n} \frac{\partial f(x^n_{n-\tau})}{\partial x_j}. \tag{23}$$

LEMMA 4.3 *Let*

(1) $\{X_j\}_{j\geq 1} \in \mathcal{S}(\alpha)$ *and* $\int_1^\infty \alpha(\tau)^\upsilon d\tau < \infty, \quad 0 < \upsilon < 1$ *hold,*
(2) $f(x^n_{n-\tau})$ *be a twice continuously differentiable function,*
(3) $b \to 0$ *and* $nb^{(\tau+1)\frac{\upsilon+1}{2}} \to \infty$ *as* $n \to \infty$.

Then the covariance of the estimate of the pdf partial derivative is bounded by

$$|C(\widehat{f}'_{x_n}(x^n_{n-\tau})| \leq$$
$$\leq \frac{R(\upsilon, x^n_{n-\tau})}{nb^{(\tau+1)\frac{\upsilon+1}{2}}} \left(b^2 V(\upsilon, x^n_{n-\tau}) + bW(\upsilon, x^n_{n-\tau}) + R(\upsilon, x^n_{n-\tau}) \right)^{1-\upsilon} \int_1^\infty \alpha(\tau)^\upsilon d\tau + o(b^2),$$

where we denote

$$V(\upsilon, x^n_{n-\tau}) = \sum_{i=n-\tau}^{n} \left(\left(\frac{(\upsilon+1)(3\upsilon-1)}{72(\upsilon-1)^3 x_n^2} + \frac{\upsilon+1}{(\upsilon-1)^2 x_i} - \frac{\upsilon(\upsilon+1)}{9(\upsilon-1)^4 x_n^3} \right) f(x^n_{n-\tau}) \right.$$
$$\left. + \frac{\upsilon+1}{\upsilon-1} \frac{\partial f(x^n_{n-\tau})}{\partial x_i} - \frac{\upsilon(\upsilon+1)}{9(\upsilon-1)^3 x_n^2} \frac{\partial f(x^n_{n-\tau})}{\partial x_n} + \frac{x_i}{2} \frac{\partial^2 f(x^n_{n-\tau})}{\partial x_i^2} \right),$$

$$W(\upsilon, x^n_{n-\tau}) = \sum_{i=n-\tau}^{n} \left(\left(\frac{3\upsilon-1}{4(\upsilon-1)x_n} + \frac{\upsilon+1}{(\upsilon-1)^2 x_i} + \frac{2x_i(\upsilon+1)}{3(\upsilon-1)^3 x_n^2} \right) f(x^n_{n-\tau}) \right.$$
$$\left. + \frac{\upsilon+1}{\upsilon-1} \frac{\partial f(x^n_{n-\tau})}{\partial x_i} + \frac{2(\upsilon+1)x_i}{3(\upsilon-1)^2 x_n} \frac{\partial f(x^n_{n-\tau})}{\partial x_n} + \frac{x_i}{2} \frac{\partial^2 f(x^n_{n-\tau})}{\partial x_i^2} \right),$$

$$L(\upsilon, x^n_{n-\tau}) = \sum_{i=n-\tau}^{n} \left(f(x^n_{n-\tau}) \frac{3\upsilon-1}{2(\upsilon-1)} + x_i \left(- \frac{4}{\upsilon-1} \frac{\partial f(x^n_{n-\tau})}{\partial x_n} - \frac{4}{(\upsilon-1)^2 x_n} f(x^n_{n-\tau}) \right) \right),$$

$$R(\upsilon, x^n_{n-\tau}) = \left(\prod_{j=n-\tau}^{n} x_j^{-\frac{\upsilon+1}{2}} \right) \frac{(2\pi)^{-\frac{(\tau+1)(\upsilon-1)}{2}-\tau}}{2x_n^2}.$$

The proofs of Lemmas 4.1 - 4.3 are given in Appendices .7, .8, .10.

Using the upper bound of the covariance $|C(\widehat{f}'_{x_n}(x^n_{n-\tau})|$ we can obtain the upper bound of the MISE and find the optimal bandwidth b.

THEOREM 4.4 *In the conditions of Lemmas 4.1 - 4.3, the optimal bandwidth that pro-*

vides minimum of the MISE is

$$b^* = \left(\frac{\tau+3}{2^\tau \pi^{\frac{\tau+1}{2}}} \frac{\int\limits_0^\infty \frac{f(x^n_{n-\tau})}{x_n} \prod\limits_{j=n-\tau}^{n} x_j^{-1/2} dx^n_{n-\tau}}{\int\limits_0^\infty \left(\frac{f(x^n_{n-\tau})}{3x_n^2} + \frac{1}{x_n} \sum\limits_{i=n-\tau}^{n} x_i \frac{\partial^2 f(x^n_{n-\tau})}{\partial x_i^2} \right)^2 dx^n_{n-\tau}} \right)^{\frac{2}{\tau+7}} n^{-\frac{2}{\tau+7}} \tag{24}$$

and the corresponding MISE has the order $MISE \sim n^{-\frac{4}{\tau+7}}$.

The proof is given in Appendix .10.

COROLLARY 4.5 *The result of Theorem 4.4 coincides with the results of (?) for univariate pdf derivative when* $\tau = 1$.

4.1. *Acknowledgements*

An unnumbered section, e.g. `\section*{Acknowledgement(s)}`, should be used for thanks, etc. and placed before any Notes or References sections.

4.2. *Funding*

An unnumbered section, e.g. `\section*{Funding}`, should be used for grant details, etc. and placed before any Notes or References sections.

.1. *Proof of Lemma 3.1*

Proof. To find the bias of the estimate $\widehat{f}(x^n_{n-\tau})$ let us write the expectation

$$\mathsf{E}_X(\widehat{f}(x^n_{n-\tau})) = \int f(t^n_{n-\tau}) \prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}(t_j)\, dt^n_{n-\tau} = \mathsf{E}_\xi(f(\xi^n_{n-\tau})), \tag{1}$$

where the r.v.s ξ_j are iid gamma distributed with the expectation $\mu_j = x_j$ and the variance $\sigma_j^2 = x_j b_j$. The product kernel $\prod\limits_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}(t_j)$ is used in (1) as a density function. To find $\mathsf{E}_\xi(f(\xi^n_{n-\tau}))$ let us do the Taylor expansion of $f(\xi^n_{n-\tau})$ in the point μ_j

$$f(\xi^n_{n-\tau}) = f(\mu^n_{n-\tau}) + \sum_{j=n-\tau}^{n} (\xi_j - \mu_j) \frac{\partial f(\xi^n_{n-\tau})}{\partial x_j}\bigg|_{\xi=\mu} + \frac{1}{2} \sum_{j=n-\tau}^{n} (\xi_j - \mu_j)^2 \frac{\partial^2 f(\xi^n_{n-\tau})}{\partial x_j^2}\bigg|_{\xi=\mu}$$
$$+ \sum_{j\neq l} (\xi_j - \mu_j)(\xi_l - \mu_l) \frac{\partial^2 f(\xi^n_{n-\tau})}{\partial x_j \partial x_l}\bigg|_{\xi=\mu} + o\left(\sum_{j=n-\tau}^{n} b_j \right).$$

Taking the expectation from the both sides of the latter equation we can write

$$\mathsf{E}_\xi(f(\xi^n_{n-\tau})) = f(x^n_{n-\tau}) + \frac{1}{2} \sum_{j=n-\tau}^{n} x_j b_j \frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2} + o\left(\sum_{j=n-\tau}^{n} b_j \right). \tag{2}$$

Hence, the bias of the multivariate pdf estimate is

$$Bias(\widehat{f}(x_{n-\tau}^n)) = \frac{1}{2}\sum_{j=n-\tau}^{n} x_j b_j \frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} + o\left(\sum_{j=n-\tau}^{n} b_j\right).$$

When $b_1 = b_2 = \ldots = b_n = b$ holds, we get (14). ■

.2. *Proof of Lemma 3.2*

Proof. By definition, the variance for the estimate $\widehat{f}(x_{n-\tau}^n)$ is

$$var(\widehat{f}(x_{n-\tau}^n)) = \frac{1}{n}\left(\mathsf{E}\left(\left(\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}(X_j)\right)^2\right) - \mathsf{E}^2\left(\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}(X_j)\right)\right). \quad (3)$$

Using the expectation (2), the second term in (3) can be written as

$$\frac{1}{n}\mathsf{E}_X^2(\widehat{f}(x_{n-\tau}^n)) = \frac{1}{n}\left(f(x_{n-\tau}^n) + \frac{1}{2}\sum_{j=n-\tau}^{n} x_j b_j \frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} + o\left(\sum_{j=n-\tau}^{n} b_j\right)\right)^2. \quad (4)$$

The first term in (3) is

$$\frac{1}{n}\mathsf{E}\left(\left(\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}(X_j)\right)^2\right) = \frac{1}{n}\int\left(\prod_{j=n-\tau}^{n} K^2_{\rho(x_j,b_j)}(X_j)\right) f(t_{n-\tau}^n)dt_{n-\tau}^n \quad (5)$$

$$= \frac{1}{n}\int\left(\prod_{j=n-\tau}^{n} \frac{t_j^{\frac{2x_j}{b_j}-2}\exp\left(\frac{-2t_j}{b_j}\right)}{b_j^{\frac{2x_j}{b_j}}\Gamma^2\left(\frac{x_j}{b_j}\right)}\right) f(t_{n-\tau}^n)dt_{n-\tau}^n = \frac{1}{n}\left(\prod_{j=n-\tau}^{n} B(x_j,b_j)\right)\mathsf{E}_\eta\left(f(\eta_{n-\tau}^n)\right),$$

where $\{\eta_j\}$are the iid gamma distributed r.v.s with the expectation $\mu_j = x_j - \frac{b_j}{2}$ and the variance $\sigma_j^2 = \frac{x_j b_j}{2} - \frac{b_j^2}{4}$. We denote

$$B(x_j,b_j) = \frac{b_j^{-3}x_j^2\Gamma(\frac{2x_j}{b_j}-1)}{2^{\frac{2x_j}{b_j}-1}\Gamma^2(\frac{x_j}{b_j}+1)}. \quad (6)$$

Let use use the notation from **?**, where $R(z) = \sqrt{2\pi}\exp(-z)z^{z+1/2}/\Gamma(z+1)$ for $z \geq 0$. Hence, we can express the gamma functions in (6) as

$$\Gamma^2\left(\frac{x}{b}+1\right) = \left(\frac{\sqrt{2\pi}\exp(\frac{-x}{b})(\frac{x}{b})^{\frac{x}{b}+1/2}}{R(\frac{x}{b})}\right)^2,$$

$$\Gamma\left(\frac{2x}{b}-1\right) = \frac{\Gamma\left(\frac{2x}{b}+1\right)}{\frac{2x}{b}\left(\frac{2x}{b}-1\right)} = \frac{\sqrt{2\pi}\exp\left(-\frac{2x}{b}\right)\left(\frac{2x}{b}\right)^{\frac{2x}{b}+\frac{1}{2}}}{\frac{2x}{b}\left(\frac{2x}{b}-1\right)R\left(\frac{2x}{b}\right)}.$$

Substituting the latter expressions in (6) we can rewrite it as

$$B(x_j, b_j) = \frac{b_j^{-\frac{1}{2}} x_j^{-\frac{1}{2}} R^2(\frac{x_j}{b_j})}{2\sqrt{\pi} R(\frac{2x_j}{b_j})(1 - \frac{b_j}{2x_j})}.$$

According to Lemma 3 in **?**, $R(z)$ is an increasing function which converges to 1 as $z \to \infty$ and $R(z) < 1$ for any $z > 0$. Then it holds

$$B(x_j, b_j) = \begin{cases} \frac{b_j^{-1/2} x_j^{-1/2}}{2\sqrt{\pi}}, & \text{if} \quad \frac{x_j}{b_j} \to \infty, \\ \frac{b^{-1} k^2 \Gamma(2k-1)}{2^{2k-1}\Gamma^2(k+1)}, & \text{if} \quad \frac{x_j}{b_j} \to k. \end{cases} \tag{7}$$

The expectation in (5) can be written similarly to (2) as

$$\mathsf{E}_\eta(f(\eta_{n-\tau}^n)) = f(\mu_{n-\tau}^n) + \frac{1}{2} \sum_{j=n-\tau}^{n} \left(\frac{x_j b_j}{2} - \frac{b_j^2}{4} \right) \frac{\partial^2 f(\mu_{n-\tau}^n)}{\partial x_j^2} + o\left(\sum_{j=n-\tau}^{n} b_j^2 \right).$$

Therefore, Taylor expanding the arguments of the functions $f(\mu_{n-\tau}^n)$

$$f(\mu_{n-\tau}^n) = f(x_{n-\tau}^n) - \sum_{j=n-\tau}^{n} \frac{b_j}{2} \frac{\partial f(x_{n-\tau}^n)}{\partial x_j} + \sum_{j=n-\tau}^{n} \frac{b_j^2}{8} \frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} + o\left(\sum_{j=n-\tau}^{n} b_j^2 \right),$$

we deduce that (5) can be written as

$$\frac{1}{n} \left(\prod_{j=n-\tau}^{n} B(x_j, b_j) \right) \left(f(x_{n-\tau}^n) - \sum_{j=n-\tau}^{n} \frac{b_j}{2} \frac{\partial f(x_{n-\tau}^n)}{\partial x_j} + \frac{b_j^2}{8} \frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} \right.$$
$$\left. + \frac{1}{2} \sum_{j=n-\tau}^{n} \left(\frac{x_j b_j}{2} - \frac{b_j^2}{4} \right) \left(\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} - \sum_{i=n-\tau}^{n} \frac{b_i}{2} \frac{\partial^3 f(x_{n-\tau}^n)}{\partial x_j^2 \partial x_i} \right) \right) + o\left(\sum_{j=n-\tau}^{n} b_j^2 \right).$$

Combining (4) and (7) we can finally write the variance of the pdf estimate as

$$var(\widehat{f}(x_{n-\tau}^n)) = \frac{1}{n} \left(\prod_{j=n-\tau}^{n} \frac{b_j^{-1/2} x_j^{-1/2}}{2\sqrt{\pi}} \right) \left(f(x_{n-\tau}^n) - \sum_{j=n-\tau}^{n} \frac{b_j}{2} \frac{\partial f(x_{n-\tau}^n)}{\partial x_j} \right.$$
$$\left. + \sum_{j=n-\tau}^{n} \frac{b_j^2}{8} \frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} + \frac{1}{2} \left(\frac{x_j b_j}{2} - \frac{b_j^2}{4} \right) \left(\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} - \sum_{i=n-\tau}^{n} \frac{b_i}{2} \frac{\partial^3 f(x_{n-\tau}^n)}{\partial x_j^2 \partial x_i} \right) \right)$$
$$- \frac{1}{n} \left(f(x_{n-\tau}^n) + \frac{1}{2} \sum_{j=n-\tau}^{n} x_j b_j \frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} \right)^2 + o\left(\sum_{j=n-\tau}^{n} b_j^2 \right).$$

When $b_1 = b_2 = \ldots = b_n = b$ holds, we get

$$var(\widehat{f}(x^n_{n-\tau})) = \frac{b^{-\frac{(\tau+1)}{2}}}{n}\left(\prod_{j=n-\tau}^{n}\frac{x_j^{-1/2}}{2\sqrt{\pi}}\right)\left(f(x^n_{n-\tau}) - \frac{b}{2}\sum_{j=n-\tau}^{n}\frac{\partial f(x^n_{n-\tau})}{\partial x_j}\right.$$
$$\left. + \frac{b^2}{8}\sum_{j=n-\tau}^{n}\frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2} + \frac{1}{2}\left(\frac{x_j b}{2} - \frac{b^2}{4}\right)\left(\frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2} - \frac{b}{2}\sum_{i=n-\tau}^{n}\frac{\partial^3 f(x^n_{n-\tau})}{\partial x_j^2 \partial x_i}\right)\right)$$
$$- \frac{1}{n}\left(f(x^n_{n-\tau}) + \frac{b}{2}\sum_{j=n-\tau}^{n} x_j \frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2}\right)^2 + o\left(b^2\right).$$

Using the notations (16) the latter variance can be written in the shorter form (15). ■

.3. *Proof of Lemma 3.3*

Proof. To evaluate the covariance of the estimate $\widehat{f}(x^n_{n-\tau})$ we shall apply Davydov's inequality (see **?**)

$$|Cov\left(\widetilde{\widetilde{K}}\left(X_1, x, b\right), \widetilde{\widetilde{K}}\left(X_{1+i}, x, b\right)\right)| \leq 2\pi\alpha(i)^{1/r} \parallel \widetilde{\widetilde{K}}\left(X_1, x, b\right) \parallel_q \parallel \widetilde{\widetilde{K}}\left(X_{1+i}, x, b\right) \parallel_p,$$

where $p^{-1} + q^{-1} + r^{-1} = 1$, $1 \leq p, q, r \leq \infty$, $\alpha(i)$ is the mixing coefficient (2). The $\mathcal{L}_p$ norm is by definition

$$\begin{aligned}
\parallel \widetilde{\widetilde{K}}\left(X_1, x, b\right) \parallel_q &= \left(\int\left(\widetilde{\widetilde{K}}\left(y, x, b\right)\right)^q f(y)dy\right)^{\frac{1}{q}} \\
&= \left(\int\left(\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}\left(t_j\right)\right)^q f(t^n_{n-\tau})dt^n_{n-\tau}\right)^{\frac{1}{q}} \\
&= \left(\mathsf{E}\left(f(\xi^n_{n-\tau})\prod_{j=n-\tau}^{n} K^{q-1}_{\rho(x_j,b_j)}\left(\xi_j\right)\right)\right)^{\frac{1}{q}}, \qquad (8)
\end{aligned}$$

where the product kernel $\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}\left(\xi_j\right)$ is used as the density function. The $\{\xi_j\}$ are the iid gamma distributed r.v.s with the expectation $\mu_j = x_j$ and the variance $var = x_j b_j$. Analogously to the previous proofs, we use the Taylor expansion to find the expectation

(8), i.e. it holds

$$\mathsf{E}\left(f(\xi_{n-\tau}^{n})\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j)}^{q-1}(\xi_j)\right)=f(\mu_{n-\tau}^{n})\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j)}^{q-1}(\mu_j) \tag{9}$$

$$+\sum_{i=n-\tau}^{n}\frac{var}{2}\frac{\partial^2}{\partial\xi_i^2}\left(f(\xi_{n-\tau}^{n})\prod_{j=n-\tau}^{n}K_{\rho_1(x_j),b_j}^{q-1}(\xi_j)\right)\Bigg|_{\xi=\mu}+o\left(\sum_{j=n-\tau}^{n}b_j^2\right)$$

$$=\left(\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j)}^{q-1}(x_j)\right)\left(f(x_{n-\tau}^{n})\frac{(3-q)}{2}+\sum_{i=n-\tau}^{n}\frac{q(q-1)b_i}{2x_i}f(x_{n-\tau}^{n})\right.$$

$$\left.-(q-1)b_i\frac{\partial f(x_{n-\tau}^{n})}{\partial x_i}+\frac{b_ix_i}{2}\frac{\partial^2 f(x_{n-\tau}^{n})}{\partial x_i^2}\right)+o\left(\sum_{j=n-\tau}^{n}b_j^2\right).$$

Using Stirling's formula

$$\Gamma(z)=\sqrt{\frac{2\pi}{z}}\left(\frac{z}{e}\right)^z\left(1+O\left(\frac{1}{z}\right)\right),$$

we can rewrite the product kernel function as

$$\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j)}(x_j)=\prod_{j=n-\tau}^{n}\frac{x_j^{\frac{x_j}{b_j}-1}\exp(-\frac{x_j}{b_j})}{b_j^{\frac{x_j}{b_j}}\Gamma(\frac{x_j}{b_j})}=\prod_{j=n-\tau}^{n}\frac{x_j^{-\frac{1}{2}}b_j^{-\frac{1}{2}}}{\sqrt{2\pi}(1+O(b_j/x_j))}.$$

Its upper bound is

$$\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j)}^{q-1}(x_j)\leq\prod_{j=n-\tau}^{n}\frac{1}{(2\pi x_jb_j)^{\frac{q-1}{2}}}. \tag{10}$$

Hence, substituting (10) in (9) we can rewrite (8) as

$$\mathsf{E}\left(f(\xi_{n-\tau}^{n})\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j)}^{q-1}(\xi_j)\right)=\left(\prod_{j=n-\tau}^{n}(2\pi x_jb_j)^{\frac{1-q}{2}}\right)\left(f(x_{n-\tau}^{n})\frac{(3-q)}{2}\right.$$

$$\left.+\sum_{i=n-\tau}^{n}\frac{q(q-1)b_i}{2x_i}f(x_{n-\tau}^{n})-(q-1)b_i\frac{\partial f(x_{n-\tau}^{n})}{\partial x_i}+\frac{b_ix_i}{2}\frac{\partial^2 f(x_{n-\tau}^{n})}{\partial x_i^2}\right)+o\left(\sum_{j=n-\tau}^{n}b_j^2\right).$$

When $b_1=b_2=\ldots=b_n=b$ holds, we get

$$\|\,\widetilde{\widetilde{K}}(X_1,x,b)\,\|_q=b^{(\tau+1)\frac{1-q}{2q}}\left(\prod_{j=n-\tau}^{n}(2\pi x_j)^{\frac{1-q}{2q}}\right)\left(f(x_{n-\tau}^{n})\frac{(3-q)}{2}\right.$$

$$\left.+b\sum_{i=n-\tau}^{n}\frac{q(q-1)}{2x_i}f(x_{n-\tau}^{n})-(q-1)\frac{\partial f(x_{n-\tau}^{n})}{\partial x_i}+\frac{x_i}{2}\frac{\partial^2 f(x_{n-\tau}^{n})}{\partial x_i^2}\right)^{1/q}\Bigg)+o\left(b^2\right).$$

Let $p = q$ and, substituting the latter norm in Davydov's inequality, we get

$$\begin{aligned}
&\left|Cov\left(\widetilde{\widetilde{K}}\left(X_1, x, b\right), \widetilde{\widetilde{K}}\left(X_{1+k}, x, b\right)\right)\right| \\
&\leq 2\pi\alpha(k)^{1/r}\left(b^{(\tau+1)\frac{1-q}{q}}\left(\prod_{j=n-\tau}^{n}(2\pi x_j)^{\frac{1-q}{q}}\right)\left(f(x_{n-\tau}^n)\frac{(3-q)}{2}\right.\right. \\
&\left.\left.+\, b\sum_{i=n-\tau}^{n}(q-1)\left(\frac{q}{2x_i}f(x_{n-\tau}^n) - \frac{\partial f(x_{n-\tau}^n)}{\partial x_i}\right) + \frac{x_i}{2}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_i^2}\right)^{\frac{2}{q}}\right)\right) + o\left(b^2\right).
\end{aligned}$$

Taking $p = q = 2 + \delta$, $r = \frac{2+\delta}{\delta}$ it can be deduced that the covariance of the pdf estimate is

$$\begin{aligned}
&\frac{2}{n}\sum_{k=1}^{n}\left(1 - \frac{k}{n}\right)Cov\left(\widetilde{\widetilde{K}}\left(X_1, x, b\right), \widetilde{\widetilde{K}}_1\left(X_{1+k}, x, b\right)\right) \\
&\leq \frac{2b^{-(\tau+1)\frac{\delta+1}{\delta+2}}}{n}(2\pi)^{1-(\tau+1)\frac{\delta+1}{\delta+2}}\left(\prod_{j=n-\tau}^{n} x_j^{-\frac{\delta+1}{\delta+2}}\right) \\
&\cdot\left(f(x_{n-\tau}^n)\frac{1-\delta}{2} + b\sum_{i=n-\tau}^{n}\frac{(\delta+1)(\delta+2)}{2x_i}f(x_{n-\tau}^n)\right. \\
&\left.- (\delta+1)\frac{\partial f(x_{n-\tau}^n)}{\partial x_i} + \frac{x_i}{2}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_i^2}\right)^{2/q}\sum_{k=1}^{n}\left(1 - \frac{k}{n}\right)\alpha(k)^{\frac{\delta}{2+\delta}} + o\left(b^2\right).
\end{aligned}$$

Let us introduce the following notations

$$S(\delta, x_{n-\tau}^n) = \sum_{i=n-\tau}^{n}\frac{(\delta+1)(\delta+2)}{2x_i}f(x_{n-\tau}^n) - (\delta+1)\frac{\partial f(x_{n-\tau}^n)}{\partial x_i} + \frac{x_i}{2}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_i^2}, \quad (11)$$

$$D(\delta, x_{n-\tau}^n) = 2(2\pi)^{1-(\tau+1)\frac{\delta+1}{\delta+2}}\left(\prod_{j=n-\tau}^{n} x_j^{-\frac{\delta+1}{\delta+2}}\right). \quad (12)$$

In these notations the covariance can be bounded by the following expression

$$\begin{aligned}
&C(\widehat{f}(x_{n-\tau}^n)) = \frac{2}{n}\sum_{i=1}^{n}\left(1 - \frac{i}{n}\right)Cov\left(\widetilde{\widetilde{K}}\left(X_1, x, b\right), \widetilde{\widetilde{K}}\left(X_{1+i}, x, b\right)\right) \\
&\leq \frac{D(\delta, x_{n-\tau}^n)}{n}b^{-(\tau+1)\frac{\delta+1}{\delta+2}}\left(f(x_{n-\tau}^n)\frac{1-\delta}{2} + bS(\delta, x_{n-\tau}^n)\right)^{\frac{2}{2+\delta}}\int_1^{\infty}\alpha(\tau)^{\frac{\delta}{2+\delta}}d\tau + o\left(b^2\right).
\end{aligned}$$

Let us denote $\frac{\delta}{2+\delta} = \upsilon$, $0 < \upsilon < 1$. Then, we get the upper bound of the covariance (17). ■

.4. *Proof of Theorem 3.4*

Proof. Combining the bias, the variance and the covariance from Lemmas 3.1 - 3.3 we can write the MSE of the pdf estimate

$$
\begin{aligned}
&MSE(\widehat{f}(x_{n-\tau}^n)) = Bias(\widehat{f}(x_{n-\tau}^n))^2 + var(\widehat{f}(x_{n-\tau}^n)) + C(\widehat{f}(x_{n-\tau}^n)) \\
&= \left(\frac{b}{2} \sum_{j=n-\tau}^{n} x_j \frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} \right)^2 - \frac{1}{n} \left(f(x_{n-\tau}^n) + \frac{b}{2} \sum_{j=n-\tau}^{n} x_j \frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2} \right)^2 \\
&+ \frac{1}{n} \left(\prod_{j=n-\tau}^{n} \frac{b^{-\frac{(\tau+1)}{2}} x_j^{-1/2}}{2\sqrt{\pi}} \right) \left(f(x_{n-\tau}^n) + b v_1(x_{n-\tau}^n) + b^2 v_2(x_{n-\tau}^n) \right) \\
&+ \frac{D(\upsilon, x_{n-\tau}^n)}{n} b^{-(\tau+1)\frac{\upsilon+1}{2}} \left(bS(\upsilon, x_{n-\tau}^n) + f(x_{n-\tau}^n) \frac{3\upsilon - 1}{2(\upsilon - 1)} \right)^{1-\upsilon} \int_1^\infty \alpha(\tau)^\upsilon d\tau + o\left(b^2\right).
\end{aligned}
$$

Since the rate of convergence in b of the covariance is larger than one for the variance when $b \to 0$ as $n \to \infty$

$$
b^{-\frac{\tau+1}{2}} < b^{-\frac{(\tau+1)(\upsilon+1)}{2}},
$$

we can neglect the variance. Taking integral from the MSE and minimizing the resulted MISE in b we get the optimal bandwidth (18). ■

.5. *Proof of Lemma 3.5*

Proof. Let us partition the covariance of the pdf estimate into two sums

$$
\begin{aligned}
C(\widehat{f}(x_{n-\tau}^n)) \leq \frac{2}{n} \sum_{\tau=2}^{c(n)} Cov \left(\widetilde{\widetilde{K}}\left(X_1, x, b\right), \widetilde{\widetilde{K}}\left(X_\tau, x, b\right) \right) \\
+ \frac{2}{n} \sum_{\tau=c(n)+1}^{\infty} Cov \left(\widetilde{\widetilde{K}}\left(X_1, x, b\right), \widetilde{\widetilde{K}}\left(X_\tau, x, b\right) \right) = I_1 + I_2.
\end{aligned} \tag{13}
$$

Using Lemma 4.3 the second sum in (13) can be bounded by

$$
I_2 \leq \frac{D(\delta, x_{n-\tau}^n)}{n} b^{-(\tau+1)\frac{\delta+1}{\delta+2}} \left(f(x_{n-\tau}^n) \frac{1-\delta}{2} + bS(\delta, x_{n-\tau}^n) \right)^{\frac{2}{2+\delta}} \sum_{\tau=c(n)}^{\infty} \alpha(\tau)^{\frac{\delta}{2+\delta}} + o\left(b^2\right),
$$

where we use the notations (11) and (12). Moreover, we can rewrite it as

$$
I_2 \leq \frac{D(\delta, x_{n-\tau}^n)}{c(n)n} b^{-(\tau+1)\frac{\delta+1}{\delta+2}} \left(f(x_{n-\tau}^n) \frac{1-\delta}{2} + bS(\delta, x_{n-\tau}^n) \right)^{\frac{2}{2+\delta}} \sum_{\tau=2}^{\infty} \tau\alpha(\tau)^{\frac{\delta}{2+\delta}} + o\left(b^2\right).
$$

Let us denote $\delta = \frac{4\kappa}{1-2\kappa}$, $0 < \kappa < \frac{1}{2}$. In this notations, we get

$$I_2 \leq \frac{D(\kappa, x_{n-\tau}^n)}{c(n)n} b^{-(\tau+1)\frac{2\kappa+1}{2}} \left(f(x_{n-\tau}^n) \frac{6\kappa - 1}{2(2\kappa - 1)} + bS(\kappa, x_{n-\tau}^n) \right)^{1-2\kappa} \sum_{\tau=2}^{\infty} \tau \alpha(\tau)^{2\kappa} + o\left(b^2\right).$$

The first sum in (13) is determined and bounded by

$$\begin{aligned}
I_1 &= \frac{2}{n} \sum_{\tau=2}^{c(n)} \left| Cov\left(\widetilde{\widetilde{K}}\left(X_1, x, b\right), \widetilde{\widetilde{K}}\left(X_\tau, x, b\right) \right) \right| \\
&= \frac{2}{n} \sum_{\tau=2}^{c(n)} \left| \mathsf{E}(\widetilde{\widetilde{K}}\left(X_1, x, b\right) \cdot \widetilde{\widetilde{K}}\left(X_\tau, x, b\right))) - \mathsf{E}(\widetilde{\widetilde{K}}\left(X_1, x, b\right)) \cdot \mathsf{E}(\widetilde{\widetilde{K}}\left(X_\tau, x, b\right)) \right| \\
&= \frac{2}{n} \sum_{\tau=2}^{c(n)} \Big| \int_0^\infty \int_0^\infty \widetilde{\widetilde{K}}\left(u, x, b\right) \widetilde{\widetilde{K}}\left(v, x, b\right) f_\tau(u, v) du dv \\
&- \int_0^\infty \widetilde{\widetilde{K}}\left(u, x, b\right) f(u) du \int_0^\infty \widetilde{\widetilde{K}}\left(v, x, b\right) f(v) dv \Big| \\
&= \frac{2}{n} \sum_{\tau=2}^{c(n)} \Big| \int_0^\infty \int_0^\infty \widetilde{\widetilde{K}}\left(u, x, b\right) \widetilde{\widetilde{K}}\left(v, x, b\right) \left(f_\tau(u, v) - f(u) f(v)\right) du dv \Big| \\
&\leq \frac{2}{n} \sum_{\tau=2}^{c(n)} \int_0^\infty \int_0^\infty \left| \widetilde{\widetilde{K}}\left(u, x, b\right) \widetilde{\widetilde{K}}\left(v, x, b\right) \right| |f_\tau(u, v) - f(u) f(v))| du dv.
\end{aligned}$$

Under the condition of the lemma $|f_\tau(x, y) - f(x) f(y)| \leq M$ we get

$$I_1 \leq \frac{2M}{n} \sum_{\tau=2}^{c(n)} \left[\int_0^\infty \Big| \prod_{j=n-\tau}^{n} K_{\rho(x_j, b_j), b_j}\left(u\right) \Big| du \right]^2 = \frac{2Mc(n)}{n}.$$

We aim to make the rate of convergence in b of I_1 larger than the rate of I_2. Then it must be

$$c(n) < b^{\frac{-(\tau+1)(2\kappa+1)}{4}}.$$

Let us choose, for example, $c(n) = b^{-\frac{\tau+1}{8}}$, $\kappa = 1/4$. Then the rates of convergence of I_1 and I_2 are given by (19). ■

.6. *Proof of Theorem 3.6*

Proof. Using the bias, the variance and the covariance from Lemmas 3.1, 3.2 and 3.5, let us write the MSE of the pdf estimate $\widehat{f}(x_{n-\tau}^n)$ and find the optimal bandwidth b that

minimizes the MISE. The MSE is

$$MSE(\widehat{f}(x^n_{n-\tau})) = \left(\frac{b}{2}\sum_{j=n-\tau}^{n} x_j \frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2}\right)^2 + \frac{1}{n}\left(b^{-\frac{(\tau+1)}{2}}\prod_{j=n-\tau}^{n}\frac{x_j^{-1/2}}{2\sqrt{\pi}}\right)$$
$$\cdot\ \left(f(x^n_{n-\tau}) + bv_1(x^n_{n-\tau}) + b^2 v_2(x^n_{n-\tau})\right) - \frac{1}{n}\left(f(x^n_{n-\tau}) + \frac{b}{2}\sum_{j=n-\tau}^{n} x_j\frac{\partial^2 f(x^n_{n-\tau})}{\partial x_j^2}\right)^2$$
$$+ \frac{2M}{nb^{\frac{\tau+1}{8}}} + \frac{D(\kappa, x^n_{n-\tau})}{nb^{\frac{(\tau+1)}{16}}}\left(f(x^n_{n-\tau})\frac{6\kappa-1}{2(2\kappa-1)} + bS(\kappa, x^n_{n-\tau})\right)^{1-2\kappa}\sum_{\tau=2}^{\infty}\tau\alpha(\tau)^{2\kappa} + o\left(b^2\right).$$

Since the rate of convergence in b of the variance is larger than the rate of the covariance we can neglect the latter. Taking integral from the MSE and minimizing it in b we find the optimal bandwidth (20). ■

.7. *Proof of Lemma 4.1*

Proof. To find the bias of the estimate $\widehat{f}'_{x_n}(x^n_{n-\tau})$ we use the same technic as in (.1). The expectation is

$$\mathsf{E}_X(\widehat{f}'_{x_n}(x^n_{n-\tau})) = \frac{1}{b_n}\mathsf{E}(\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}(X_j)\, L(X_n, x_n, b_n))$$
$$= \frac{1}{b_n}\int L(t_n, x_n, b_n) f(t^n_{n-\tau})\prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}(t_j)\, dt^n_{n-\tau} = \frac{1}{b_n}\mathsf{E}_\xi(f(\xi^n_{n-\tau})L(\xi_n, x_n, b_n))$$
$$= \frac{1}{b_n}\mathsf{E}_\xi(f(\xi^n_{n-\tau})\ln(\xi_n)) - \frac{1}{b_n}\mathsf{E}_\xi(f(\xi^n_{n-\tau})(\ln b_n + \Psi(\rho(x_n, b_n)))) + o\left(\sum_{j=n-\tau}^{n} b_j\right), \quad (14)$$

where $\{\xi_j\}$ are the iid gamma distributed r.v.s with the expectation $\mu_j = x_j$ and the variance $x_j b_j$. Using the Taylor expansion in $\mu^n_{n-\tau}$, we can write

$$f(\xi^n_{n-\tau})\ln(\xi_n) = f(\mu^n_{n-\tau})\ln(\mu_n) + \sum_{j=n-\tau}^{n}(\xi_j - \mu_j)\frac{\partial\left(f(\xi^n_{n-\tau})\ln(\xi_n)\right)}{\partial x_j}\Big|_\mu$$
$$+ \frac{1}{2}\sum_{j=n-\tau}^{n}(\xi_j-\mu_j)^2\frac{\partial^2\left(f(\xi^n_{n-\tau})\ln(\xi_n)\right)}{\partial x_j^2}\Big|_\mu$$
$$+ \sum_{j\neq l}(\xi_j-\mu_j)(\xi_l-\mu_l)\frac{\partial^2\left(f(\xi^n_{n-\tau})\ln(\xi_n)\right))}{\partial x_j\partial x_l}\Big|_\mu + o\left(\sum_{j=n-\tau}^{n} b_j\right).$$

Taking the expectation from the right and the left sides of the latter we get the first term in (14)

$$\mathsf{E}_{\xi}(f(\xi_{n-\tau}^{n})\ln(\xi_{n})) = f(x_{n-\tau}^{n})\ln(x_{n}) + \frac{1}{2}\sum_{j=n-\tau}^{n} x_{j}b_{j}\frac{\partial^{2}f}{\partial x_{j}^{2}}\ln(x_{n}) + \frac{1}{2}x_{n}b_{n}\left(\frac{2}{x_{n}}\frac{\partial f}{\partial x_{n}} - \frac{f(x_{n-\tau}^{n})}{x_{n}^{2}}\right) + o\left(\sum_{j=n-\tau}^{n} b_{j}\right).$$

For the second term in (14) we use the approximation of the Digamma function when $\rho \to \infty$

$$\Psi(\rho) = \ln\rho - \frac{1}{2\rho} - \frac{1}{12\rho^{2}} + \frac{1}{120\rho^{4}} - \frac{1}{252\rho^{6}} + O\left(\frac{1}{\rho^{8}}\right).$$

Hence, we can write that

$$\ln b_{n} + \Psi(\rho(x_{n}, b_{n})) = \ln x_{n} - \frac{b_{n}}{2x_{n}} - \frac{b_{n}^{2}}{12x_{n}^{2}} + o(b_{n}^{2}).$$

Hence (14) can be rewritten as follows

$$\mathsf{E}_{X}(\widehat{f}'_{x_{n}}(x_{n-\tau}^{n})) = f(x_{n-\tau}^{n})\frac{b_{n}}{12x_{n}^{2}} + \frac{\partial f(x_{n-\tau}^{n})}{\partial x_{n}} + \frac{1}{2}\sum_{j=n-\tau}^{n} x_{j}b_{j}\frac{\partial^{2}f(x_{n-\tau}^{n})}{\partial x_{j}^{2}}\left(\frac{1}{2x_{n}} + \frac{b_{n}}{12x_{n}^{2}}\right) + o\left(\sum_{j=n-\tau}^{n} b_{j}\right). \quad (15)$$

When $\tau = 0$ holds, the multivariate expectation (15) coincides with its univariate version in **?**. From (15) we get the bias of the pdf derivative estimate

$$Bias(\widehat{f}'_{x_{n}}(x_{n-\tau}^{n})) = f(x_{n-\tau}^{n})\frac{b_{n}}{12x_{n}^{2}} + \frac{1}{2}\sum_{j=n-\tau}^{n} x_{j}b_{j}\frac{\partial^{2}f(x_{n-\tau}^{n})}{\partial x_{j}^{2}}\left(\frac{1}{2x_{n}} + \frac{b_{n}}{12x_{n}^{2}}\right) + o\left(\sum_{j=n-\tau}^{n} b_{j}\right).$$

When $b_{1} = b_{2} = \ldots = b_{n} = b$ holds, it follows that the Bias of the density derivative is (21). ■

.8. *Proof of Lemma 4.2*

Proof. By definition, the variance for the estimate $\widehat{f}'_{x_n}(x^n_{n-\tau})$ is

$$var(\widehat{f}'_{x_n}(x^n_{n-\tau})) = \frac{1}{n} Var\left(\frac{1}{b_n} L(X_n, x_n, b_n) \prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}(X_j)\right)$$

$$= \frac{1}{nb_n^2}\mathsf{E}\left(L^2(X_n, x_n, b_n) \prod_{j=n-\tau}^{n} K^2_{\rho(x_j,b_j)}(X_j)\right)$$

$$- \frac{1}{nb_n^2}\mathsf{E}^2\left(L(X_n, x_n, b_n) \prod_{j=n-\tau}^{n} K_{\rho(x_j,b_j)}(X_j)\right). \tag{16}$$

The second term of the right-hand side of (16) is the square degree of (14). Hence using (15) we can immediately write that

$$\frac{1}{n}\mathsf{E}^2_X(\widehat{f}'_{x_n}(x^n_{n-\tau})) = \frac{1}{n}\Bigg(Bias^2(\widehat{f}'_{x_n}(x^n_{n-\tau})) + \left(\frac{\partial f(x^n_{n-\tau})}{\partial x_n}\right)^2$$

$$+ 2\frac{\partial f(x^n_{n-\tau})}{\partial x_n} Bias(\widehat{f}'_{x_n}(x^n_{n-\tau})) + o\left(\sum_{j=n-\tau}^{n} b_j^2\right)\Bigg).$$

The first term in (16) can be represented as

$$\frac{1}{nb_n^2}\mathsf{E}\left(L^2(X_n, x_n, b_n) \prod_{j=n-\tau}^{n} K^2_{\rho(x_j,b_j)}(X_j)\right)$$

$$= \frac{1}{nb_n^2}\int L^2(t_n, x_n, b_n) f(t^n_{n-\tau}) \prod_{j=n-\tau}^{n} K^2_{\rho(x_j,b_j)}(t_j)\, dt^n_{n-\tau}$$

$$= \frac{1}{nb_n^2}\int \left(\prod_{j=n-\tau}^{n} \frac{t_j^{\frac{2x_j}{b_j}-2}\exp\left(\frac{-2t_j}{b_j}\right)}{b_j^{\frac{2x_j}{b_j}}\Gamma^2\left(\frac{x_j}{b_j}\right)}\right) L^2(t_n, x_n, b_n) f(t^n_{n-\tau}) dt^n_{n-\tau}. \tag{17}$$

Using the property of the gamma function $\Gamma^2(\frac{x}{b}+1)=(\frac{x}{b})^2\Gamma^2(\frac{x}{b})$, (17) can be rewritten as

$$\frac{1}{nb_n^2}\int\left(\prod_{j=n-\tau}^{n}\frac{b_j^{-3}x_j^2\Gamma(\frac{2x_j}{b_j}-1)}{2^{\frac{2x_j}{b_j}-1}\Gamma^2(\frac{x_j}{b_j}+1)}\frac{t_j^{\frac{2x_j}{b_j}-2}\exp(\frac{-2t_j}{b_j})}{\left(\frac{b_j}{2}\right)^{\frac{2x_j}{b_j}-1}\Gamma(\frac{2x_j}{b_j}-1)}\right)L^2(t_n,x_n,b_n)f(t_{n-\tau}^n)dt_{n-\tau}^n$$

$$=\frac{1}{nb_n^2}\left(\prod_{j=n-\tau}^{n}B(x_j,b_j)\right)\mathsf{E}_\eta\left(L^2(\eta_n,x_n,b_n)f(\eta_{n-\tau}^n)\right)$$

$$=\frac{1}{nb_n^2}\left(\prod_{j=n-\tau}^{n}B(x_j,b_j)\right)\left(\mathsf{E}_\eta\left(f(\eta_{n-\tau}^n)\ln^2\eta_n\right)+\left(\ln b_n+\Psi\left(\frac{x_n}{b_n}\right)\right)^2\mathsf{E}_\eta\left(f(\eta_{n-\tau}^n)\right)\right.$$

$$\left.-2\left(\ln b_n+\Psi\left(\frac{x_n}{b_n}\right)\right)\mathsf{E}_\eta\left(f(\eta_{n-\tau}^n)\ln\eta_n\right)\right), \tag{18}$$

where $\{\eta_j\}$ are the iid gamma distributed r.v.s with the expectation $\mu_j=x_j-\frac{b_j}{2}$ and the variance $\frac{x_jb_j}{2}-\frac{b_j^2}{4}$, and $B(x_j,b_j)$ is determined by (7). To determine the expectations in (18) we use the same technique as before. Namely, the expectations are

$$\mathsf{E}_\eta(f(\eta_{n-\tau}^n)\ln(\eta_n))=f(\mu_{n-\tau}^n)\left(\ln\mu_n-\frac{b_n}{4\mu_n}\right)+\sum_{j=n-\tau}^{n}\left(\frac{x_jb_j}{2}-\frac{b_j^2}{4}\right)\frac{\partial^2f(\mu_{n-\tau}^n)}{\partial x_j^2}\frac{\ln\mu_n}{2}$$

$$+\frac{b_n}{2}\frac{\partial f(\mu_{n-\tau}^n)}{\partial x_n}+o\left(\sum_{j=n-\tau}^{n}b_j^2\right),$$

$$\mathsf{E}_\eta(f(\eta_{n-\tau}^n))=f(\mu_{n-\tau}^n)+\frac{1}{2}\sum_{j=n-\tau}^{n}\left(\frac{x_jb_j}{2}-\frac{b_j^2}{4}\right)\frac{\partial^2f(\mu_{n-\tau}^n)}{\partial x_j^2}+o\left(\sum_{j=n-\tau}^{n}b_j^2\right).$$

$$\mathsf{E}_\eta(f(\eta_{n-\tau}^n)\ln^2(\eta_n))=f(\mu_{n-\tau}^n)\left(\ln^2(\mu_n)+\frac{b_n}{2\mu_n}(1-\ln\mu_n)\right)$$

$$+\frac{\ln^2\mu_n}{2}\sum_{j=n-\tau}^{n}\left(\frac{x_jb_j}{2}-\frac{b_j^2}{4}\right)\frac{\partial^2f(\mu_{n-\tau}^n)}{\partial x_j^2}+b_n\ln\mu_n\frac{\partial f(\mu_{n-\tau}^n)}{\partial x_n}+o\left(\sum_{j=n-\tau}^{n}b_j^2\right).$$

Using the known Tailor expansions

$$\ln\mu_n=\ln x_n-\frac{b_n}{2x_n}-\frac{b_n^2}{8x_n^2}+o(b_n^2),$$

$$\frac{1}{\mu_n}=\frac{1}{x_n}+\frac{b_n}{2x_n^2}+\frac{b_n^2}{4x_n^3}+o(b_n^2)$$

we can rewrite (18) as follows

$$\frac{1}{n}\left(\prod_{j=n-\tau}^{n}B(x_j,b_j)\right)\left(f(\mu_{n-\tau}^n)\left(\frac{1}{4x_n^2}+\frac{1}{2x_nb_n}+\frac{7b_n}{48x_n^3}+\frac{7b_n^2}{576x_n^4}+\frac{b_n^3}{192x_n^5}\right)\right.$$
$$\left.+\sum_{j=n-\tau}^{n}\left(\frac{x_jb_j}{2}-\frac{b_j^2}{4}\right)\frac{\partial^2 f(\mu_{n-\tau}^n)}{\partial x_j^2}\frac{b_n^2}{1152x_n^4}-\frac{b_n}{24x_n^2}\frac{\partial f(\mu_{n-\tau}^n)}{\partial x_n}+o\left(\sum_{j=n-\tau}^{n}b_j^2\right)\right).$$

Therefore, Taylor expanding the arguments of the functions $f(\mu_{n-\tau}^n)$ in $x_{n-\tau}^n$, substituting it in latter and using (17), we can write the variance of the pdf derivative estimate as

$$var(\widehat{f}'_{x_n}(x_{n-\tau}^n))=$$
$$=\frac{1}{n}\left(\prod_{j=n-\tau}^{n}\frac{b_j^{-1/2}x_j^{-1/2}}{2\sqrt{\pi}}\right)\left(-\frac{b_n}{24x_n^2}\left(\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}-\frac{b_j}{2}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_n\partial x_j}+\sum_{j=n-\tau}^{n}\frac{b_j^2}{4}\frac{\partial^3 f(x_{n-\tau}^n)}{\partial x_n\partial x_j^2}\right)\right.$$
$$+\frac{b_n^2}{1152x_n^4}\sum_{j,i=n-\tau}^{n}\left(\frac{x_jb_j}{2}-\frac{b_j^2}{4}\right)\left(\frac{b_i^2}{4}\frac{\partial^4 f(x_{n-\tau}^n)}{\partial x_j^2\partial x_i^2}-\frac{b_i}{2}\frac{\partial^3 f(x_{n-\tau}^n)}{\partial x_j^2\partial x_i}\right)$$
$$+\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2}\left(\left(\frac{x_jb_j}{2}-\frac{b_j^2}{4}\right)\frac{b_n^2}{1152x_n^4}+\frac{b_j^2}{4}\left(\frac{1}{4x_n^2}+\frac{1}{2x_nb_n}+\frac{7b_n}{48x_n^3}+\frac{7b_n^2}{576x_n^4}+\frac{b_n^3}{192x_n^5}\right)\right)$$
$$+\left(f(x_{n-\tau}^n)-\frac{b_j}{2}\frac{\partial f(x_{n-\tau}^n)}{\partial x_j}\right)\left(\frac{1}{4x_n^2}+\frac{1}{2x_nb_n}+\frac{7b_n}{48x_n^3}+\frac{7b_n^2}{576x_n^4}+\frac{b_n^3}{192x_n^5}\right)$$
$$\left.-\frac{1}{n}\left(Bias^2(\widehat{f}'_{x_n}(x_{n-\tau}^n))+\left(\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}\right)^2+2\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}Bias(\widehat{f}'_{x_n}(x_{n-\tau}^n))\right)+o\left(\sum_{j=n-\tau}^{n}b_j^2\right)\right).$$

If $b_1=b_2=\ldots=b_n=b$ hold, we get

$$var(\widehat{f}'_{x_n}(x_{n-\tau}^n))=$$
$$=\frac{b^{-\frac{(\tau+1)}{2}}}{n}\left(\prod_{j=n-\tau}^{n}\frac{x_j^{-1/2}}{2\sqrt{\pi}}\right)\left(-\frac{b}{24x_n^2}\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}+\sum_{j=n-\tau}^{n}\frac{b^2}{48x_n^2}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_n\partial x_j}\right.$$
$$+\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_j^2}\left(\frac{b^2}{16x_n^2}+\frac{b}{8x_n}\right)+f(x_{n-\tau}^n)\left(\frac{1}{4x_n^2}+\frac{1}{2x_nb}+\frac{7b}{48x_n^3}+\frac{7b^2}{576x_n^4}\right)$$
$$-\frac{\partial f(x_{n-\tau}^n)}{\partial x_j}\left(\frac{b}{8x_n^2}+\frac{1}{4x_n}+\frac{7b^2}{96x_n^3}\right)$$
$$\left.-\frac{1}{n}\left(Bias^2(\widehat{f}'_{x_n}(x_{n-\tau}^n))+\left(\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}\right)^2+2\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}Bias(\widehat{f}'_{x_n}(x_{n-\tau}^n))\right)+o(b^2).\right.$$

Introducing the notations (23), we can write the variance of the pdf derivative estimate in form (22). ■

.9. *Proof of Lemma 4.3*

Proof. Furthermore, we apply Davydov's inequality for the covariance of the pdf derivative estimate

$$|Cov\left(\widetilde{K}\left(X_1,x,b\right),\widetilde{K}\left(X_{1+i},x,b\right)\right)|\leq 2\pi\alpha(i)^{1/r}\parallel\widetilde{K}\left(X_1,x,b\right)\parallel_q\parallel\widetilde{K}\left(X_{1+i},x,b\right)\parallel_p,$$

where $p^{-1}+q^{-1}+r^{-1}=1$, $1\leq p,q,r\leq\infty$, $\alpha(i)$ is the mixing coefficient (2). The $\mathcal{L}_p$ norm is by definition

$$\begin{aligned}
&\parallel\widetilde{K}\left(X_1,x,b\right)\parallel_q=\left(\int\left(\widetilde{K}\left(y,x,b\right)\right)^q f(y)dy\right)^{1/q}\\
&=\left(\int\left(L(t_n,x_n,b_n)\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j),b_j}\left(t_j\right)\right)^q f(t_{n-\tau}^n)dt_{n-\tau}^n\right)^{1/q}\\
&=\left(\int\left(L^q(t_n,x_n,b_n)f(t_{n-\tau}^n)\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j),b_j}^{q-1}\left(t_j\right)\right)\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j),b_j}\left(t_j\right)dt_{n-\tau}^n\right)^{1/q}\\
&=\left(\mathsf{E}\left(L(\xi_n,x_n,b_n)^q f(\xi_{n-\tau}^n)\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j),b_j}^{q-1}\left(\xi_j\right)\right)\right)^{1/q},
\end{aligned}\tag{19}$$

where the kernel $\prod_{j=n-\tau}^{n}K_{\rho_1(x_j),b_j}\left(\xi_j\right)$ is used as the density function and $\{\xi_j\}$ are the iid gamma distributed r.v.s with the expectation $\mu_j=x_j$ and the variance x_jb_j. The Taylor expansion of (19) is the following

$$\begin{aligned}
&L(\xi_n,x_n,b_n)^q f(\xi_{n-\tau}^n)\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j),b_j}^{q-1}\left(\xi_j\right)\\
&=L(\mu_n,x_n,b_n)^q f(\mu_{n-\tau}^n)\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j),b_j}^{q-1}\left(\mu_j\right)\\
&+\sum_{i=n-\tau}^{n}\left(\xi_i-\mu_i\right)\frac{\partial}{\partial\xi_i}\left(L(\xi_n,x_n,b_n)^q f(\xi_{n-\tau}^n)\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j),b_j}^{q-1}\left(\xi_j\right)\right)\Bigg|_{\xi=\mu}\\
&+\sum_{i\neq l}\frac{(\xi_i-\mu_i)(\xi_l-\mu_l)}{2}\frac{\partial^2}{\partial\xi_i^2}\left(L(\xi_n,x_n,b_n)^q f(\xi_{n-\tau}^n)\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j),b_j}^{q-1}\left(\xi_j\right)\right)\Bigg|_{\xi=\mu}\\
&+\sum_{i=n-\tau}^{n}\frac{(\xi_i-\mu_i)^2}{2}\frac{\partial^2}{\partial\xi_i^2}\left(L(\xi_n,x_n,b_n)^q f(\xi_{n-\tau}^n)\prod_{j=n-\tau}^{n}K_{\rho(x_j,b_j),b_j}^{q-1}\left(\xi_j\right)\right)\Bigg|_{\xi=\mu}+o\left(\sum_{j=n-\tau}^{n}b_j^2\right).
\end{aligned}$$

Taking the expectation from the latter expression and using the known Taylor expansions

$$L(x_n,x_n,b_n)=\ln x_n-\ln b_n-\Psi\left(\frac{x_n}{b_n}\right)=\frac{b_n}{2x_n}+\frac{b_n^2}{12x_n^2}+o(b_n^2),\tag{20}$$

$$L^q(x_n, x_n, b_n) = \left(\frac{b_n}{2x_n}\right)^q \left(1 + q\frac{b_n}{6x_n} + \frac{q(q-1)}{2}\left(\frac{b_n}{6x_n}\right)^2\right),$$

$$\frac{\partial}{\partial \xi_n} L(\xi_n, x_n, b_n) = \frac{1}{\xi_n}\Big|_{\xi_n = x_n} = \frac{1}{x_n}, \quad \frac{\partial^2}{\partial \xi_n^2} L(\xi_n, x_n, b_n) = -\frac{1}{\xi_n}\Big|_{\xi_n = x_n} = -\frac{1}{x_n^2},$$

the expectation (19) can be rewritten as

$$\begin{aligned}
&\mathsf{E}\left(\prod_{j=n-\tau}^{n} K^{q-1}_{\rho_1(x_j),b_j}(\xi_j)\, L_1(\xi_n, x_n, b_n)^q f(\xi^n_{n-\tau})\right) \\
&= \left(\prod_{j=n-\tau}^{n} K^{q-1}_{\rho_1(x_j),b_j}(x_j)\right)\left(\left(\frac{b_n}{2x_n}\right)^q \left(1 + q\frac{b_n}{6x_n} + \frac{q(q-1)}{2}\left(\frac{b_n}{6x_n}\right)^2\right) f(x^n_{n-\tau})\right. \\
&+ \left(\frac{b_n}{2x_n}\right)^{q-1} \sum_{i=n-\tau}^{n} \frac{x_i b_i}{2}\left(\left(\frac{b_n}{2x_n} + \frac{qb_n^2}{12x_n^2} + \frac{q(q-1)b_n^3}{144x_n^3}\right)\left(f(x^n_{n-\tau})\left(\frac{1-q}{b_i x_i} - \frac{q(1-q)}{x_i^2}\right)\right.\right. \\
&+ \left.\frac{\partial f(x^n_{n-\tau})}{\partial x_i}\frac{2(1-q)}{x_i} + \frac{\partial^2 f(x^n_{n-\tau})}{\partial x_i^2}\right) - \left(1 + \frac{(q-1)b_n}{6x_n} + \frac{(q-1)(q-2)b_n^2}{72x_n^2}\right) \\
&\left.\left. \cdot \left(f(x^n_{n-\tau})\frac{q^2}{x_n^2} - \frac{2q}{x_n}\frac{\partial f(x^n_{n-\tau})}{\partial x_n}\right)\right) + o\left(\sum_{j=n-\tau}^{n} b_j^2\right).
\end{aligned}$$

If $b_1 = b_2 = \ldots = b_n = b$ hold and, using (10), we can obtain that

$$\begin{aligned}
&\| \widetilde{K}(X_1, x, b) \|_q \\
&\le \left(\prod_{j=n-\tau}^{n} (2\pi x_j)^{\frac{1-q}{2q}}\right) \frac{b^{(\tau+1)\frac{1-q}{2q}+1}}{2x_n}\left(\sum_{i=n-\tau}^{n}\left(1 + \frac{qb}{6x_n} + \frac{q(q-1)b^2}{72x_n^2}\right)\right. \\
&\cdot \left(f(x^n_{n-\tau})\frac{3-q}{2} + b(1-q)\frac{\partial f(x^n_{n-\tau})}{\partial x_i} + \frac{x_i b}{2}\frac{\partial^2 f(x^n_{n-\tau})}{\partial x_i^2} - \frac{q(1-q)b}{2x_i} f(x^n_{n-\tau})\right) \\
&\left. + x_i\left(1 + \frac{(q-1)b}{6x_n} + \frac{(q-1)(q-2)b^2}{72x_n^2}\right)\left(2q\frac{\partial f(x^n_{n-\tau})}{\partial x_n} - \frac{q^2}{x_n} f(x^n_{n-\tau})\right)\right)^{1/q} + o\left(b^2\right).
\end{aligned}$$

The same is valid for $\| \widetilde{K}(X_{1+i}, x, b) \|_p$. Let us take $p = q$. Hence, the covariance of the pdf derivative estimate is

$$
\begin{aligned}
&|Cov\left(\widetilde{K}(X_1, x, b), \widetilde{K}(X_{1+k}, x, b)\right)| \\
&\leq 2\pi\alpha(k)^{1/r}\Bigg(\left(\prod_{j=n-\tau}^{n}(2\pi x_j)^{\frac{1-q}{q}}\right)\frac{b^{(\tau+1)\frac{1-q}{q}+2}}{4x_n^2}\Bigg(\sum_{i=n-\tau}^{n}\left(1+\frac{qb}{6x_n}+\frac{q(q-1)b^2}{72x_n^2}\right) \\
&\cdot\left(f(x_{n-\tau}^n)\frac{3-q}{2}+b(1-q)\frac{\partial f(x_{n-\tau}^n)}{\partial x_i}+\frac{x_i b}{2}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_i^2}-\frac{q(1-q)b}{2x_i}f(x_{n-\tau}^n)\right) \\
&+x_i\left(1+\frac{(q-1)b}{6x_n}+\frac{(q-1)(q-2)b^2}{72x_n^2}\right)\left(2q\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}-\frac{q^2}{x_n}f(x_{n-\tau}^n)\right)\Bigg)^{2/q}+o\left(b^2\right).
\end{aligned}
$$

Taking $p = q = 2+\delta$, $r = \frac{2+\delta}{\delta}$ and using the following notations

$$
\begin{aligned}
V(\delta, x_{n-\tau}^n) = \sum_{i=n-\tau}^{n}\Bigg(&-(\delta+1)\frac{\partial f(x_{n-\tau}^n)}{\partial x_i}+\frac{\delta(\delta+1)(\delta+2)}{36x_n^2}\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}+\frac{x_i}{2}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_i^2} \\
&+\left(-\frac{(\delta-1)(\delta+1)(\delta+2)}{144x_n^2}+\frac{(\delta+1)(\delta+2)}{2x_i}-\frac{\delta(\delta+1)(\delta+2)^2}{72x_n^3}\right)f(x_{n-\tau}^n)\Bigg),
\end{aligned}
$$

$$
\begin{aligned}
W(\delta, x_{n-\tau}^n) = \sum_{i=n-\tau}^{n}\Bigg(&\left(\frac{1-\delta}{4x_n}+\frac{(\delta+1)(\delta+2)}{2x_i}-\frac{x_i(\delta+1)(\delta+2)^2}{6x_n^2}\right)f(x_{n-\tau}^n) \\
&-(\delta+1)\frac{\partial f(x_{n-\tau}^n)}{\partial x_i}+\frac{(\delta+1)(\delta+2)x_i}{3x_n}\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}+\frac{x_i}{2}\frac{\partial^2 f(x_{n-\tau}^n)}{\partial x_i^2}\Bigg),
\end{aligned}
$$

$$
L(\delta, x_{n-\tau}^n) = \sum_{i=n-\tau}^{n}\left(f(x_{n-\tau}^n)\frac{(1-\delta)}{2}+x_i\left(2(2+\delta)\frac{\partial f(x_{n-\tau}^n)}{\partial x_n}-\frac{(2+\delta)^2}{x_n}f(x_{n-\tau}^n)\right)\right),
$$

$$
R(\delta, x_{n-\tau}^n) = \left(\prod_{j=n-\tau}^{n}x_j^{-\frac{\delta+1}{\delta+2}}\right)\frac{(2\pi)^{\frac{\tau+1}{\delta+2}-\tau}}{2x_n^2},
$$

the covariance of the pdf derivative estimate is

$$
\begin{aligned}
&C(\widehat{f}'_{x_n}(x_{n-\tau}^n) = \frac{2}{nb^2}\sum_{k=1}^{n}\left(1-\frac{k}{n}\right)Cov\left(\widetilde{K}(X_1, x, b), \widetilde{K}(X_{1+k}, x, b)\right) \\
&\leq \frac{R(\delta, x_{n-\tau}^n)}{b^{(\tau+1)\frac{\delta+1}{\delta+2}}n}\left(b^2V(\delta, x_{n-\tau}^n)+bW(\delta, x_{n-\tau}^n)+L(\delta, x_{n-\tau}^n)\right)^{2/(\delta+2)}\int_1^{\infty}\alpha(\tau)^{\frac{\delta}{2+\delta}}d\tau+o\left(b^2\right).
\end{aligned}
$$

Let us denote $\frac{\delta}{2+\delta} = \upsilon$, $0 < \upsilon < 1$. Then, in this notations, we get (24). ■

.10. *Proof of Theorem 4.4*

Proof. Using the bias, the variance and the covariance in Lemmas 4.1, 4.2, 4.3, we can write the MSE for the estimate of the pdf derivative

$$
\begin{aligned}
&MSE(\widehat{f}'_{x_n}(x^n_{n-\tau})) = (Bias\widehat{f}'_{x_n}(x^n_{n-\tau}))^2 + var(\widehat{f}'_{x_n}(x^n_{n-\tau})) + C(\widehat{f}'_{x_n}(x^n_{n-\tau})) \\
&= \left(bB_1(x^n_{n-\tau}) + b^2B_2(x^n_{n-\tau})\right)^2 \left(1 - \frac{1}{n}\right) \\
&+ \frac{b^{-\frac{(\tau+1)}{2}}}{n}\left(\prod_{j=n-\tau}^{n}\frac{x_j^{-1/2}}{2\sqrt{\pi}}\right)\left(bV_1(x^n_{n-\tau}) + b^2V_2(x^n_{n-\tau}) + \frac{1}{b}V_3(x^n_{n-\tau}) + V_4(x^n_{n-\tau})\right) \\
&- \frac{1}{n}\left(\left(\frac{\partial f(x^n_{n-\tau})}{\partial x_n}\right)^2 + 2\frac{\partial f(x^n_{n-\tau})}{\partial x_n}\left(bB_1(x^n_{n-\tau}) + b^2B_2(x^n_{n-\tau})\right)\right) + \frac{R(\upsilon, x^n_{n-\tau})}{n}b^{-(\tau+1)\frac{\upsilon+1}{2}} \\
&\cdot \left(b^2V(\upsilon, x^n_{n-\tau}) + bW(\upsilon, x^n_{n-\tau}) + L(\upsilon, x^n_{n-\tau})\right)^{1-\upsilon}\int_1^\infty \alpha(\tau)^\upsilon d\tau + o\left(b^2\right).
\end{aligned}
$$

Since the rate of convergence of the covariance is less than one of the variance, we can neglect the covariance term. Taking the integral from the MSE and minimizing the MISE in b the optimal bandwidth (24) is obtained. ■